



Pediatric Tachycardia – With Pulses

Approval: Troy M. Falck, MD – Medical Director

Effective: 12/01/2020

Approval: Victoria Pinette – Executive Director

Next Review: 11/2023

Signs of Cardiopulmonary Compromise

- Acutely altered mental status

- Hypotension

- Signs of shock

BLS

- Assess & support ABC's
- High flow O₂
- Assess V/S, including SpO₂

ALS

- Cardiac monitor, 12-lead ECG at appropriate time
- Consider IV/IO NS (may bolus 20 mL/kg)

Probable Sinus Tachycardia

- P waves present & normal
- Variable R-R & constant P-R
- Infants: rate usually <220
- Children: rate usually <180

Treat underlying cause

Monitor & reassess

Probable SVT

- P waves absent or abnormal
- HR not variable
- Infants: rate usually ≥220
- Children: rate usually ≥180

Cardiopulmonary compromise?

NO

Vagal Maneuver

Vagal Maneuver successful?

NO

Adenosine (Base Hospital Order Only)

- 1st dose: 0.1 mg/kg rapid IV/IO (max 6 mg), followed with 20 mL IV/IO NS flush
- If rhythm does not convert within 1-2 min:
- 2nd dose: 0.2 mg/kg rapid IV/IO (max 12 mg), followed with 20 mL IV/IO NS flush

Probable VT

Cardiopulmonary compromise?

NO

YES

Synchronized Cardioversion
Base Hospital Order Only

- Initial dose: 0.5-1 J/kg
- Subsequent doses: 2 J/kg
- Consider sedation

Contact base hospital for consultation as needed