



**Acute Respiratory Distress**

Approval: Troy M. Falck, MD – Medical Director

Effective: 12/01/2020

Approval: Victoria Pinette – Executive Director

Next Review: 11/2023

**Continuous Positive Airway Pressure (CPAP) Utilization**

**• Indications:**

- CHF with pulmonary edema
- Moderate to severe respiratory distress
- Near drowning

**• Contraindications:**


- <8 years of age
- Respiratory or cardiac arrest
- Severe decreased LOC
- Agonal respirations
- Inability to maintain airway
- Suspected pneumothorax
- SBP <90
- Major trauma, especially head injury or significant chest trauma

**• Complications:**

- Hypotension
- Pneumothorax
- Corneal drying

**Epinephrine Administration**

- Epinephrine is only indicated for pts with suspected asthma who are in severe distress.
- **Use epinephrine cautiously in pts >35yo, or with a history of coronary artery disease or hypertension.**
- Administer Auto-Injector/IM epinephrine into the lateral thigh, midway between waist & knee.



**SEE PAGE 2 FOR ALS TREATMENT**



### Acute Respiratory Distress

#### Asthma/COPD

#### ALS

##### Mild Distress

- Mild wheezing
- Mild shortness of breath
- Cough

- Cardiac monitor
- Consider IV NS (may bolus up to 1000 mL)

- Albuterol 5 mg & Ipratropium 500 mcg**
- Nebulizer
- May repeat (**albuterol 2.5-5 mg only**) for continued respiratory distress

##### Moderate to Severe Distress

- Cyanosis
- Accessory muscle use
- Inability to speak >3 words
- Severe wheezing/shortness of breath
- Decreased or absent air movement

- Cardiac monitor
- IV/IO NS (may bolus up to 1000 mL)

- Albuterol 5 mg & Ipratropium 500 mcg**
- Nebulizer/CPAP/BVM
- May repeat (**albuterol 2.5-5 mg only**) for continued respiratory distress

##### History of asthma with severe distress only

- Epinephrine 1:1,000**
- 0.01 mg/kg IM (max: 0.5 mg)

#### CHF/Pulmonary Edema

#### ALS

##### Mild Signs & Symptoms

- Cardiac monitor
- IV NS TKO

- \*Nitroglycerin**
- 0.4 mg SL
- May repeat every 5 mins

##### **\*Nitroglycerin Notes/Precautions**

- Do not administer if SBP <100
- Do not delay due to difficult vascular access
- Consult with base/modified base hospital prior to administration to pts taking erectile dysfunction or pulmonary HTN medication

##### Moderate to Severe Signs & Symptoms

- Cardiac monitor
- Assess BP x 2 to confirm accuracy
- IV/IO NS TKO

- \*Nitroglycerin: titrate dose based on SBP**
- SBP 100-150: 0.4 mg SL
- SBP 150-200: 0.8 mg SL
- SBP >200: 1.2 mg SL
- May repeat titrated doses every 5 mins based on repeat SBP