

Sierra – Sacramento Valley EMS Agency Program Policy

Needle Cricothyrotomy

	Effective: 12/01/2021	Next Review: 09/2024	1103
	Approval: Troy M. Falck, MD – Medical Director		SIGNATURE ON FILE
	Approval: Victoria Pinette – Executive Director		SIGNATURE ON FILE

PURPOSE:

To establish needle cricothyrotomy criteria for S-SV EMS accredited paramedics.

AUTHORITY:

- A. HSC § 1797.220 and 1798.
- B. CCR, Title 22, Division 9, Chapter 4.

POLICY:

- A. Needle cricothyrotomy may be performed by paramedic personnel, when there is an inability to maintain the airway utilizing less invasive airway procedures. This situation typically involves patients with pathologic processes that cause distortion of the upper airway anatomy, including one or more of the following:
 - 1. Airway obstruction by uncontrolled bleeding into the oral cavity and/or vomiting.
 - 2. Severe maxillofacial trauma - blunt, penetrating, or associated with mandibular FX.
 - 3. Laryngeal foreign body that cannot be removed expeditiously.
 - 4. Swelling of upper airway structures.
 - 5. Infection (e.g., epiglottitis, Ludwig's angina).
 - 6. Allergic reaction or hereditary angioedema.
 - 7. Chemical or thermal burns to the epiglottis and upper airway.
- B. Needle cricothyrotomy is contraindicated in the following circumstances:
 - 1. Patient age <3 years or estimated weight <15 kg.
 - 2. Ability to maintain airway utilizing less invasive procedures.
 - 3. Conscious patient.
 - 4. Moving ambulance.
 - 5. Patient has a midline neck hematoma or massive subcutaneous emphysema.