

Sierra – Sacramento Valley EMS Agency Treatment Protocol

Suspected Moderate/Severe Traumatic Brain Injury (TBI)

Effective: 04/01/2025

T-3

Approval: Troy M. Falck, MD – Medical Director

Approval: John Poland – Executive Director

Next Review: 01/2028

Prehospital Identification of Moderate/Severe TBI

- Any pt with a mechanism of injury consistent with a potential for a brain injury, and one or more of the following:
 - <65 years of age with a GCS ≤13, or ≥65 years of age with a GCS <15 (or decrease from baseline)
 - Post-traumatic seizures
 - Multi-system trauma requiring advanced airway placement

For any patient with a suspected moderate/severe TBI, avoid/treat the three TBI "H-Bombs":

1) Hyperventilation, 2) Hypoxia, 3) Hypotension



- Assess V/S, including continuous SpO₂ monitoring and pupil exam: Reassess V/S every 3-5 min if possible
- High-flow O₂ (regardless of SpO₂ reading)
- If continued hypoxia (SpO₂ <94%) or inadequate ventilatory effort, proceed through the following in a stepwise manner
 - Reposition airway
 - Initiate positive pressure ventilation with appropriate airway adjunct if necessary (use of a pressurecontrolled BVM &/or ventilation rate timer is recommended if available)
- Avoid hyperventilation (ventilate at a rate of 10 breaths/min)

Monitor & reassess

- Maintain normothermia
- Consider the concurrent need for appropriate immobilization/spinal motion restriction



- Continuous cardiac & EtCO2 monitoring
- Check blood glucose

