



**Pediatric Respiratory Distress**

Approval: Troy M. Falck, MD – Medical Director

Effective: 04/01/2026

Approval: John Poland – Executive Director

Next Review: 01/2029

- Consider respiratory failure for pts with a history of increased work of breathing & presenting with ALOC & a slow or normal respiratory rate without retractions.
- The hallmark of upper airway obstruction (croup, epiglottitis, foreign body airway obstruction) is inspiratory stridor.
- Do not attempt to visualize the throat or insert anything into the mouth if epiglottitis suspected.

**Continuous Positive Airway Pressure (CPAP) Utilization Information**

**• Indications:**

- CHF with pulmonary edema
- Moderate to severe respiratory distress
- Near drowning

**• Contraindications:**

- <8 years of age
- Agonal respirations
- SBP <90
- Respiratory or cardiac arrest
- Inability to maintain airway
- Major trauma/head injury/chest trauma
- Suspected croup/epiglottitis
- Suspected pneumothorax
- Severe decreased LOC

**• Complications:**

- Hypotension
- Pneumothorax
- Corneal drying

**Epinephrine Administration**

- Epinephrine is indicated for pts with suspected asthma who are in severe distress.
- Administer Auto-Injector/IM epinephrine into the lateral thigh, midway between waist & knee.

**BLS**

- Assess & support ABCs
- High flow O<sub>2</sub>
- Assess V/S, including SpO<sub>2</sub>
- Assess history & physical, determine degree of illness
- Minimize stimulation – keep pt calm & consider allowing parent to hold the child &/or O<sub>2</sub> delivery device if their presence calms the child
- Consider CPAP, when appropriate/indicated, for moderate to severe distress (pts ≥8 yo only)

**Suspected asthma & in severe distress**

**YES**

**Epinephrine 1:1,000 IM (authorized/trained EMTs only)**

- Pts 7.5 – 30 kg
  - 0.15 mg pediatric auto-injector **OR** 0.15 mg (0.15 mL) via approved syringe
- Pts >30 kg
  - 0.3 mg adult auto-injector **OR** 0.3 mg (0.3 mL) via approved syringe

**See page 2 for LALS treatment of wheezing or suspected croup/epiglottitis**



**Pediatric Respiratory Distress**

Wheezing

LALS

**Mild Distress**

- Mild wheezing
- Mild shortness of breath
- Cough

- Cardiac & EtCO<sub>2</sub> monitoring (**AEMT II**)

**Albuterol 5 mg**

- Nebulizer
- May repeat 2.5-5 mg for continued respiratory distress

**Moderate – Severe Distress**

- Cyanosis
- Accessory muscle use
- Inability to speak >3 words
- Severe wheezing/shortness of breath
- Decreased or absent air movement

- Cardiac & EtCO<sub>2</sub> monitoring (**AEMT II**)
- IV/IO NS (may bolus 20 mL/kg)

**Albuterol 5 mg**

- Nebulizer, CPAP, or BVM
- May repeat 2.5-5 mg for continued respiratory distress

**Epinephrine 1:1,000 (for severe distress only)**

- 0.01 mg/kg IM (max: 0.3 mg)

Suspected Croup/Epiglottitis

LALS

- Cardiac & EtCO<sub>2</sub> monitoring (**AEMT II**)
- Consider nebulized saline
- **Minimize pt stimulation**
- If full upper airway occlusion suspected – ensure proper airway positioning & BVM seal, attempt to ventilate & reassess – **do not attempt i-gel**

**Base/Modified Base Hospital Order Only**

**Racemic epinephrine**

- One (1) - 0.5 ml vial of 2.25% inhalation solution (mix with NS to = 5 ml of volume)

**OR**

**Nebulized epinephrine**

- 1:1000 - 0.5 mL/kg (max: 5 mL) nebulizer or BVM (if <5 mL, mix with NS to = 5 mL of volume)